arXiv:1303.3143v1 [gr-qc] 13 Mar 2013

On Quasinormal Modes for Gravitational Perturbations of Regular Black Holes

S. C. Ulhoa*
Faculdade Gama, Universidade de Brasília, 72444-240, Setor Leste (Gama), Brasília, DF, Brazil.

In this paper we investigate gravitational perturbations of regular black holes. Such systems are solutions of Einstein equations that do not have a singularity at the origin of the coordinate system. However they still have events horizons depending on the values of the characteristic parameters of the solutions. When a black hole is perturbed, it oscillates. It gives rise to damped vibrating modes which are known as quasinormal modes. It is calculated the quasinormal frequencies of regular black holes using the third order WKB approximation for gravitational perturbations. The results are presented in tables I, II, III and IV.

PACS numbers: 04.20-q; 04.20.Cv; 02.20.Sv
Keywords: Quasinormal frequencies; Gravitational perturbations; Regular black holes; Bardeen spacetime.

I. INTRODUCTION

In 1957 Regge and Wheeler explored the stability of Schwarzschild solution over gravitational perturbations [1]. They discovered that it leads to a Schröedinger-like equation with a specific effective potential and a frequency associated to the temporal dependence. Since then others perturbations for different systems have been analyzed and lead to similar equations [2]. The frequencies in those equations can be calculated by means semi-analytic or numerical methods. Such frequencies are discrete and complex numbers which can be obtained by the imposition of specific boundary conditions such as the existence of ongoing waves at the spatial infinity and ingoing waves at the event horizon . This is known as quasinormal modes. These modes are damped oscillations of the spacetime geometry that can be used to investigate fundamental features of the gravitational field. The real part is related to the oscillation itself while the imaginary part is refers to the rate at each mode is damped. Therefore quasinormal modes may be important to gravitational waves physics and it could become an environment to study basic effects of quantum gravity since the quasinormal frequencies are discrete quantities.

In what concerns black holes the presence of a fundamental singularity can cause serious difficulties in making predictions which lead Einstein himself to doubt about the physical realization of the first solutions of the field equations of general relativity [3]. Over the years a great effort has been directed to deal with such problem by means the development of the so called cosmic censorship hypotheses [4, 5]. The singularity would be inaccessible to an observer outside the event horizon and always covered by it which allows the avoidance of a naked singularity. This is a route to contour the problem, not to solve it, since the singularity, which is a place where no physics can be done, is still there. Perhaps this might be solved by the introduction of regular black holes. Such objects are interesting because they do not have a fundamental singularity at the origin of the coordinate system. Event horizons can be present but they could be removed under some change of variables.

The first regular solution of Einstein equations that describes a black hole was due to Bardeen who got his solution only approximatively [6]. Later it was discovered that such a solution could be viewed as the gravitational collapse of a magnetic monopole to which a nonlinear electromagnetic energy-momentum tensor works as the source of field equations which were exactly solved [7]. Others regular solutions was obtained by the coupling of some matter field, such as scalar field, to gravitation in a cosmological context [8]. Because of their singularity-free feature regular black holes could be great candidates to represent realistic final stages of collapsing regular configurations such as a common star. Although the success of the computation of quasinormal modes of black holes or neutron stars, it has not been done for regular black holes for gravitational perturbations. This is exactly what it is intended to be studied in this paper.

The paper is organized as follows. In section II the general theory of black holes stability of a spherical symmetric background metric for gravitational perturbations is summarized. The master equation and the effective potential are obtained. Then three different solutions are introduced, among then the Bardeen solution which was the first one describing a regular black hole. In section III the complex frequencies for each solution has been calculated. It was done using WKB approximation of third order, the results are then organized in tables I, II, II and IV. Finally in the last section the concluding remarks are presented.

II. GRAVITATIONAL PERTURBATIONS OF REGULAR BLACK HOLES

In this section we carry out a treatment to deal with a gravitational perturbation of regular black holes. A generic spherically symmetric background metric $\eta_{\mu\nu}$, is defined by the line element

---

* sc.ulhoa@gmail.com

$$ds^2 = -f(r)\,dt^2 + f^{-1}(r)\,dr^2 + r^2(d\theta^2 + sin^2\theta d\phi^2)\,, \quad (1)$$

thus the metric tensor can be written as $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$. The perturbations $h_{\mu\nu}$ can be expanded in terms of the spherical harmonics which, in the Regge-Wheeler gauge [1], reads

$$h_{\mu\nu} = \begin{pmatrix} 0 & 0 & 0 & h_0(r,t) \\ 0 & 0 & 0 & h_1(r,t) \\ 0 & 0 & 0 & 0 \\ h_0(r,t) & h_1(r,t) & 0 & 0 \end{pmatrix} \sin\theta\,\partial_\theta P_l(\theta)\,, \quad (2)$$

where $P_l(\theta)$ are the Legendre polynomials.

The perturbed Einstein equations are given by

$$\delta R_{\mu\nu} = -\partial_\alpha \delta\Gamma^\alpha_{\mu\nu} + \partial_\nu \delta\Gamma^\alpha_{\mu\alpha}\,,$$

where

$$\delta\Gamma^\alpha_{\beta\gamma} = \frac{1}{2}\eta^{\alpha\nu}(\partial_\gamma h_{\beta\nu} + \partial_\beta h_{\gamma\nu} - \partial_\nu h_{\beta\gamma})\,.$$

Thus, from eq. (2), the above equations yields

$$\frac{1}{f(r)}\frac{\partial h_0}{\partial t} - \frac{\partial[f(r)h_1]}{\partial r} = 0\,, \quad (3)$$

$$\frac{1}{f(r)}\left[\frac{\partial^2 h_1}{\partial t^2} - \frac{\partial^2 h_0}{\partial t \partial r} + \frac{2}{r}\frac{\partial h_0}{\partial t}\right] + \frac{1}{r^2}[l(l+1) - 2]h_1 = 0\,, \quad (4)$$

for the functions $h_0(r,t)$ and $h_1(r,t)$.

If we use the definition $\psi = \left(\frac{1}{r}\right) f(r) h_1(r,t)$, then eqs. (3) and (4) can be combined into a single one which reads

$$\frac{\partial^2\psi}{\partial t^2} - f^2\frac{\partial^2\psi}{\partial r^2} - f\frac{\partial f}{\partial r}\frac{\partial\psi}{\partial r} + \frac{f}{r}\left[\frac{l(l+1) + 2(f-1)}{r} - \frac{\partial f}{\partial r}\right]\psi = 0\,.$$

In order to put such equation in a more familiar form, we change the variable r to the "tortoise" coordinate, defined by $dx = \frac{dr}{f(r)}$, that leads to

$$\frac{\partial^2\psi}{\partial t^2} - \frac{\partial^2\psi}{\partial x^2} + V(x)\,\psi = 0\,, \quad (5)$$

with the effective potential given by

$$V = \frac{f}{r}\left[\frac{l(l+1) + 2(f-1)}{r^2} - \frac{\partial f}{\partial r}\right]\,.$$

The temporal dependence is assumed to be given by $\psi = e^{-\imath\omega t}\phi$, thus we have

$$\left[\frac{\partial^2}{\partial x^2} + \omega^2 - V(x)\right]\phi(x) = 0\,, \quad (6)$$

which is the master equation for gravitational perturbations of a spherically symmetric background metric. The choice of $f(r)$ determines which spacetime we are dealing with. In the next subsections we introduce different solutions which are viewed as regular black holes.

A. Bardeen Space-Time

The Bardeen spacetime is settled by means the choice

$$f(r) = 1 - \frac{2mr^2}{(r^2 + \alpha^2)^{3/2}}\,,$$

where α represents a magnetic monopole and m its mass. It was shown that Bardeen black hole can be obtained as an exact solution of Einstein equations coupled to a model of nonlinear electrodynamics [6]. In other words it can be viewed as a system composed by a self-gravitating magnetic monopole of charge α and mass m. Such solution was first obtained only approximately and it was the first regular black hole ever proposed.

An interesting feature of the Bardeen solution is that it can have zero, one or two horizons of events depending on the choice of the magnetic charge α. Although it is always a regular black hole at $r = 0$, it describes a regular spacetime only when the following inequality holds:

$$\alpha^2 \leq \frac{16}{27}m^2\,.$$

There are lots of works in the literature concerning Bardeen black hole. Geodesic structure of test particles was studied in [9], while the features of Bardeen solution as a gravitational lens was given in [10]. The quasinormal modes of such a spacetime, for scalar field perturbations, was calculated in [11]. Here we intend to calculate the quasinormal frequencies for gravitational perturbations.

B. Solution II: Hayward solution

Similar to Bardeen solution, the choice

$$f(r) = 1 - \frac{2mr^2}{(r^3 + 2L^2\,m)}$$

also describes a regular black hole [12]. It is obtained in the context of nonlinear electrodynamics coupled to gravity, where the parameter L is a fundamental length scale related to the magnetic charge. Here we explore the quasinomal modes for gravitational perturbations for a specific choice of L.

C. Solution III: coupling gravity to a phantom scalar field

The following solution

$$f(r) = 1 + \frac{cr^2}{b^2} + \frac{\rho_0 r^2}{b^3}\left[\frac{b\sqrt{r^2 - b^2}}{r^2} + \tan^{-1}\left(\frac{\sqrt{r^2 - b^2}}{b}\right)\right]$$

is obtained in a cosmological context, when a phantom scalar field is coupled to Einstein equations, as an attempt to explain observational data concerning dark energy. For details about the constants of the solution see ref. [8]. For certain range of the above parameters such a solution describes a regular black hole. Indeed c, b and ρ_0 have to be positive numbers.

III. QUASINORMAL MODES IN WKB APPROXIMATION

The WKB method is a semi-analytic technique used to solve Schöedinger-type equations like eq. (6). Once the solution $\phi(x)$ is known it is possible to use two conditions to properly establish it: it is requested that there are only ongoing waves at the infinity and only ingoing waves at the horizon of events. Usually the frequency in WKB approximation is written as [13]

$$\frac{\imath\left(\omega^2-V_0\right)}{\sqrt{-2V_0''}}-\sum_{i=2}^{k}\Lambda_i=n+\frac{1}{2}\,,$$

where V_0 and V_0'' are the effective potential and its second derivative respectively, taken at the point of the maximum of V. The second term in the left-hand side of the above equation represents the higher order WKB corrections (from 2nd until the k-th order). In general a great feature of the WKB method is that it gets better and better as one take the higher orders of approximation (for low values of n and l). As a matter of fact it is a very accurate procedure even when compared to numeric methods [13].

In this paper we will work with WKB approximation of third order [14] which is given by

$$\omega_{n,l}^2=\left[V_0+(-2V_0'')^{1/2}\Lambda\right]-\imath\left(n+\frac{1}{2}\right)(-2V_0'')^{1/2}(1+\Omega)\,, \tag{7}$$

where

$$\Lambda=\frac{1}{(-2V_0'')^{1/2}}\left[\frac{1}{8}\left(\frac{V_0^{(4)}}{V_0''}\right)\left(\frac{1}{4}+\beta^2\right)-\frac{1}{288}\left(\frac{V_0'''}{V_0''}\right)^2\left(7+60\beta^2\right)\right]\,,$$

and

$$\begin{aligned}\Omega=\frac{1}{(-2V_0'')}&\left[\frac{5}{6912}\left(\frac{V_0'''}{V_0''}\right)^4\left(77+188\beta^2\right)-\frac{1}{384}\left(\frac{(V_0''')^2V_0^{(4)}}{(V_0'')^3}\right)\left(51+100\beta^2\right)+\right.\\&\left.+\frac{1}{2304}\left(\frac{V_0^{(4)}}{V_0''}\right)^2\left(67+68\beta^2\right)+\frac{1}{288}\left(\frac{V_0'''V_0^{(5)}}{(V_0'')^2}\right)\left(19+28\beta^2\right)-\frac{1}{288}\left(\frac{V_0^{(6)}}{V_0''}\right)\left(5+4\beta^2\right)\right],\end{aligned}$$

Here $\beta=n+\frac{1}{2}$ and $V_0^{(n)}=\frac{d^nV}{dx^n}|_{x=x_0}$. The point x_0 is the solution of the equation $\frac{dV}{dx}(x_0)=0$, which means that it is a point of maximum of the effective potential. It should be noted that x is the well known tortoise coordinate.

Using expression (7) it is possible to calculate the quasinormal frequencies which are presented in tables I and II for Bardeen space-time. In tables III and IV it is showed the results for solutions of subsections II B and II C respectively which were obtained with the help of the same expression for $\omega_{n,l}$. All results refer to quasinormal modes for gravitational perturbations.

We see that the quasinormal modes of Bardeen spacetime for $\alpha=\frac{4}{\sqrt{27}}$ are very similar to the results obtained for the Schwarzschild spacetime which is very unusual since such a solution is obtained from Bardeen spacetime by means $\alpha=0$. Thus it would be expected that the quasinormal modes would approximate the results of Schwarzschild spacetime for values of α close to zero.

The results obtained for Hayward solution are charted on table I. We see that the imaginary part of the frequency has a little variation which means an approximately constant damping of the quasinormal modes. On the other hand, we see that the results on table IV represent highly damped modes and a rapid oscillation as well. Thus a regular black hole described by the function $f(r)$, defined in subsection II C, seems to reach a stable configuration faster than the others solutions.

IV. CONCLUSION

In this article we have analyzed gravitational perturbations of regular black holes, by calculating the quasinormal frequencies. Such results were obtained for three solutions of Einstein equations that represent regular black

TABLE I. Quasinormal modes of Bardeen space-time for $\alpha = 0.3$. It was calculated in units of m.

l	n	$\omega_{n,l}$	l	n	$\omega_{n,l}$
2	0	$0.14845 - 0.10716i$	4	0	$0.46625 - 0.11159i$
	1	$0.13011 - 0.33531i$		1	$0.45724 - 0.33840i$
	2	$0.10465 - 0.56443i$		2	$0.44492 - 0.56935i$
	3	$0.06807 - 0.79400i$		3	$0.43043 - 0.80164i$
				4	$0.41309 - 1.03418i$
3	0	$0.32484 - 0.11046i$	5	0	$0.59773 - 0.11210i$
	1	$0.31297 - 0.33768i$		1	$0.59043 - 0.33866i$
	2	$0.29809 - 0.56899i$		2	$0.57958 - 0.56886i$
	3	$0.27943 - 0.80073i$		3	$0.56696 - 0.80095i$
	4	$0.25585 - 1.03254i$		4	$0.55244 - 1.03366i$

TABLE II. Quasinormal modes of Bardeen space-time for $\alpha = \frac{4}{\sqrt{27}}$. It was calculated in units of m.

l	n	$\omega_{n,l}$	l	n	$\omega_{n,l}$
2	0	$0.26655 - 0.07346i$	4	0	$0.63345 - 0.07782i$
	1	$0.23183 - 0.22989i$		1	$0.61816 - 0.23526i$
	2	$0.17674 - 0.39981i$		2	$0.58964 - 0.39719i$
	3	$0.10519 - 0.57786i$		3	$0.55055 - 0.56432i$
				4	$0.50267 - 0.73603i$
3	0	$0.46091 - 0.07687i$	5	0	$0.79835 - 0.07824i$
	1	$0.44009 - 0.23398i$		1	$0.78618 - 0.23585i$
	2	$0.40304 - 0.39837i$		2	$0.76293 - 0.39649i$
	3	$0.35392 - 0.56948i$		3	$0.73028 - 0.56108i$
	4	$0.29413 - 0.74562i$		4	$0.68972 - 0.72965i$

holes. To obtain the quasinormal modes it has been used third order WKB approximation which yields good results when compared to accurate numerical technics, for low values of the numbers n and l.

The results are summarized in tables I, II, III and IV. It is interesting to note that for Bardeen space-time it is much closer to Schwarzschild space-time for bigger values of α than for values of such parameter closer to zero. The quasinormal modes for Schwarzschild were already known, for instance see table III of reference [14], thus such modes are not charted here. For $\alpha = 0$ Bardeen space-time is formally equal to Schwarzschild space-time, therefore it is such a surprise that the maximum allowed value of α approximate Bardeen and Schwarzschild solutions in what concerns quasinormal modes for gravitational perturbations.

TABLE III. Quasinormal modes for Bardeen-like solution of subsection II B, using $L = 0.5$.

l	n	$\omega_{n,l}$	l	n	$\omega_{n,l}$
2	0	$0.14746 - 0.10502i$	4	0	$0.46201 - 0.10995i$
	1	$0.12987 - 0.32875i$		1	$0.45274 - 0.33341i$
	2	$0.10593 - 0.55343i$		2	$0.43986 - 0.56104i$
	3	$0.07161 - 0.77844i$		3	$0.42455 - 0.79006i$
				4	$0.40612 - 1.01936i$
3	0	$0.32184 - 0.10869i$	5	0	$0.59234 - 0.11050i$
	1	$0.30962 - 0.33243i$		1	$0.58486 - 0.33380i$
	2	$0.29412 - 0.56034i$		2	$0.57358 - 0.56071i$
	3	$0.27461 - 0.78874i$		3	$0.56026 - 0.78954i$
	4	$0.24993 - 1.01726i$		4	$0.54481 - 1.01901i$

TABLE IV. Quasinormal modes for solution of subsection II C, using $c = \rho_0 = 1$ and $b = 0.2$.

l	n	$\omega_{n,l}$	l	n	$\omega_{n,l}$
2	0	$75.47946 - 18.07363i$	4	0	$105.22227 - 20.80790i$
	1	$73.11341 - 48.44850i$		1	$105.02646 - 62.19756i$
	2	$57.34869 - 70.95716i$		2	$104.26975 - 103.26486i$
	3	$19.20005 - 96.07928i$		3	$102.80119 - 144.18648i$
				4	$100.63482 - 185.08368i$
3	0	$88.72793 - 19.50189i$	5	0	$123.45462 - 21.93977i$
	1	$87.46532 - 56.71597i$		1	$124.15338 - 65.81798i$
	2	$81.88287 - 91.59152i$		2	$125.54218 - 109.69983i$
	3	$70.59940 - 125.87692i$		3	$127.61642 - 153.59710i$
	4	$53.73096 - 161.19276i$		4	$130.37542 - 197.52066i$

---

[1] Tullio Regge and John A. Wheeler. Stability of a schwarzschild singularity. *Phys. Rev.*, 108:1063–1069, Nov 1957.

[2] Kostas D. Kokkotas and Bernd Schmidt. Quasi-normal modes of stars and black holes. *Living Reviews in Relativity*, 2(2), 1999.

[3] A. Einstein. On a stationary system with spherical symmetry consisting of many gravitating masses. *Annals of Mathematics*, 40:922–936, 1939.

[4] R. Penrose. *Riv. Nuovo Cimento*, 1:252, 1969.

[5] Sanjay M. Wagh. Classical formulation of cosmic censorship hypothesis. 2002.

[6] *Proceedings of GR5, Tbilisi, USSR, p. 174*, 1968.

[7] Eloy Ayón-Beato and Alberto García. Regular black hole in general relativity coupled to nonlinear electrodynamics. *Phys. Rev. Lett.*, 80:5056–5059, Jun 1998.

[8] K.A. Bronnikov and J.C. Fabris. Regular phantom black holes. *Phys.Rev.Lett.*, 96:251101, 2006.

[9] Sheng Zhou, Juhua Chen, and Yongjiu Wang. Geodesic Structure of Test Particle in Bardeen Spacetime. *Int.J.Mod.Phys.*, D21:1250077, 2012.

[10] Ernesto F Eiroa and Carlos M Sendra. Gravitational lensing by a regular black hole. *Classical and Quantum Gravity*, 28(8):085008, 2011.

[11] Sharmanthie Fernando and Juan Correa. Quasinormal Modes of Bardeen Black Hole: Scalar Perturbations. *Phys.Rev.*, D86:064039, 2012.

[12] Sean A. Hayward. Formation and evaporation of regular black holes. *Phys.Rev.Lett.*, 96:031103, 2006.

[13] R.A. Konoplya and A. Zhidenko. Quasinormal modes of black holes: From astrophysics to string theory. *Rev.Mod.Phys.*, 83:793–836, 2011.

[14] Sai Iyer. Black-hole normal modes: A wkb approach. ii. schwarzschild black holes. *Phys. Rev. D*, 35:3632–3636, Jun 1987.